

Use this calendar to keep track of your NINLARO® (ixazomib) regimen.


























NINLARO is a prescription medicine used to treat multiple myeloma in combination with the medicines REVLIMID® (lenalidomide) and dexamethasone, in people who have received at least one prior treatment for their multiple myeloma.

It is not known if NINLARO is safe and effective in children.

NINLARO is a capsule that you take once a week for 3 weeks of a 4-week cycle. NINLARO is taken along with 2 other medications, lenalidomide and dexamethasone, in 4-week cycles. This dosing calendar is a snapshot of the NINLARO regimen for a 28-day cycle.

See the next page for more information on how to take NINLARO.

Start by filling in the dates from Day 1 up to Day 28. Then after you take your medication, check off the corresponding box to remember what you've taken that day.

 NINLARO		 Lenalidomide		 Dexamethasone		
Date: Day 1	Date: Day 2	Date: Day 3	Date: Day 4	Date: Day 5	Date: Day 6	Date: Day 7
 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>
Date: Day 8	Date: Day 9	Date: Day 10	Date: Day 11	Date: Day 12	Date: Day 13	Date: Day 14
 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>
Date: Day 15	Date: Day 16	Date: Day 17	Date: Day 18	Date: Day 19	Date: Day 20	Date: Day 21
 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>	 <input type="checkbox"/>
Date: Day 22	Date: Day 23	Date: Day 24	Date: Day 25	Date: Day 26	Date: Day 27	Date: Day 28
 <input type="checkbox"/>	Days 23-28: No dosing					

Please read the Important Safety Information on the next page and Patient Information in the accompanying full Prescribing Information.

How should I take NINLARO® (ixazomib)?

- **Take NINLARO, REVLIMID® (lenalidomide) and dexamethasone** exactly as your healthcare provider tells you to take it
- **Take each dose of NINLARO** at about the same time of day each week
- **Take each dose of NINLARO** at least 1 hour before or at least 2 hours after food (swallow NINLARO whole with water)
- **Do not take NINLARO** at the same time as dexamethasone

Please read the [Patient Information](#) in the full [Prescribing Information](#) for more details on how to take NINLARO

Important Safety Information for NINLARO

NINLARO may cause serious side effects, including:

- **Low platelet counts (thrombocytopenia)** are common with NINLARO and can sometimes be serious. You may need platelet transfusions if your counts are too low. Tell your healthcare provider if you have any signs of low platelet counts, including bleeding and easy bruising.
- **Stomach and intestinal (gastrointestinal) problems.** Diarrhea, constipation, nausea, and vomiting are common with NINLARO and can sometimes be severe. Call your healthcare provider if you get any of these symptoms and they do not go away during treatment with NINLARO. Your healthcare provider may prescribe medicine to help treat your symptoms.
- **Nerve problems** are common with NINLARO and may also be severe. Tell your healthcare provider if you get any new or worsening symptoms including: tingling, numbness, pain, a burning feeling in your feet or hands, or weakness in your arms or legs.
- **Swelling** is common with NINLARO and can sometimes be severe. Tell your healthcare provider if you develop swelling in your arms, hands, legs, ankles, or feet, or if you gain weight from swelling.
- **Skin Reactions.** Tell your healthcare provider if you get a new or worsening rash.
- **Liver problems.** Tell your healthcare provider if you get these signs of a liver problem: yellowing of your skin or the whites of your eyes; pain in your right upper-stomach area.

Other common side effects have occurred. Tell your healthcare provider if you get new or worsening back pain, lowered white blood cells (neutropenia) that may increase the risk of infection, or vision conditions such as blurred vision, dry eye, or pink eye (conjunctivitis).

These are not all the possible side effects of NINLARO. Talk to your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before taking NINLARO, tell your healthcare provider about all your medical conditions, including if:

- You have liver problems or kidney problems or are on dialysis.
- You or your partner are pregnant or plan to become pregnant. NINLARO can harm your unborn baby. Avoid becoming pregnant during treatment with NINLARO. You and your partner should use effective birth control during treatment and for 90 days after the final dose of NINLARO. If using hormonal contraceptives (for example, the pill), an additional barrier method of contraception (for example, diaphragm or condom) must be used.
- You are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with NINLARO and for 90 days after your final dose of NINLARO.

Tell your healthcare provider about all the medications (prescription and over-the-counter) and nutritional supplements you are taking or before starting any new medicines.

Please read the [Patient Information](#) in the accompanying full [Prescribing Information](#).



ONCOLOGY

All trademarks are the property of their respective owners.

©2018 Millennium Pharmaceuticals, Inc., a wholly owned subsidiary of Takeda Pharmaceutical Company Limited.
All rights reserved. Printed in USA 8/18 MAT-US-IXA-18-00320